You notice it in the mirror first, usually while brushing. The gumline on a canine looks a little higher than its neighbor. A bit more tooth showing, a sliver of yellow that was not there last winter. Then comes the zing of cold water. Welcome to gum recession, the quiet creep that turns daily routines into a science experiment. If you walk into a Winnipeg dental office with that complaint, the evaluation you get is more than a quick glance and a "brush softer." A careful dentist in Winnipeg treats gum recession like a detective case, and the evidence starts before you even open wide.

What counts as recession, and why the millimeters matter

Gum recession is the apical migration of the gingival margin, which is a technical way of saying the pink stuff slid down the root. Teeth are not meant to have exposed roots. Dentin, which lives under the enamel and covers the root, is porous and sensitive. Roots also pick up plaque more easily and develop cavities faster than enamel-covered surfaces. Minor recession can be a cosmetic nuisance; moderate to severe recession can mean cold sensitivity, notched toothbrush abrasion, increased risk of root caries, and eventual tooth mobility.

In the chair, a winnipeg dentist measures recession in millimeters, site by site. A single millimeter may not sound dramatic, but two additional millimeters over a few years tells a different story. Rate of change is as important as the snapshot. A small, stable recession defect with thick, healthy tissue might only need monitoring and habit changes. A growing defect on a thin, delicate tissue biotype calls for a closer look and often a referral.

The interview that sets the stage

Before anyone touches a probe, the dentist asks questions. This is where the story takes shape.

- What do you feel, and when did you first notice it?
- Any cold sensitivity, sharp brushing pain, or bleeding?
- How do you brush and how often? Manual or powered brush, soft or medium bristles, pressure level?
- Flossing or interdental brushes, and how enthusiastically do you use them?
- Any clenching or grinding, especially during stressful seasons or on night shifts?
- Orthodontic history, such as previous crowding corrected by expansion or proclination?
- Past gum disease treatment, scaling visits, or deep cleanings?
- Systemic factors, including diabetes control, autoimmune disease, recent hormonal shifts, and medications like isotretinoin or calcium channel blockers?
- Tobacco, vaping, or oral piercings?
- Sports and instruments, for example hockey mouthguards or a clarinet embouchure that rubs the lower incisors?

Winnipeg winters do not cause recession, but habits that come with long cold months can nudge sensitive teeth. People clench more when they are tense. Dry air and mouth breathing from a stubborn January cold can irritate the gingiva. A dentist winnipeg patients trust asks about seasons and routine changes, because gum tissue tells time in more than months.

First look: tissue tone, biotype, and the lay of the land

Once the story is on record, the visual exam begins. Healthy gingiva has a matte surface, coral pink with stippling in some areas. Recession is clear when the yellow root shows, but there are subtler signs: rolled margins, shiny inflamed tissue that bleeds at a look, or a blanched line where a tight frenum tugs the gum every time you smile.

Biotype matters. Thin, delicate gums, the kind where you can almost see a hint of the root underneath, are beautiful and temperamental. They recede faster and graft less predictably without careful planning. Thick, fibrotic tissue is tougher and more forgiving. The width of keratinized tissue is measured too. If the band is narrow or absent on a facial surface, the gum is more vulnerable to brushing trauma and muscle pull.

The dentist also scans for toothbrush abrasion notches, V-shaped grooves right at the gumline, and for abfraction lesions, which are stress-related wedge defects. It is often a mix: a bit of aggressive brushing plus a thin biotype plus nighttime clenching. No single villain, more like a committee.

Probing, charting, and numbers with meaning

A periodontal probe is a millimeter ruler on a stick, and it drives this part of the evaluation. The dentist or hygienist charts six sites per tooth: mesio-buccal, mid-buccal, disto-buccal, mesio-lingual, mid-lingual, and disto-lingual. Two core measurements matter.

- Probing depths: the distance from the gum margin to where the probe meets tissue resistance. Healthy sulci rarely exceed 3 millimeters, though 4s can appear in tight spaces on big molars.
- Recession: the distance from the cementoenamel junction, the CEJ, to the gum margin. Positive numbers indicate recession; a negative number means the margin is coronal to the CEJ.

Add those together, and you get clinical attachment level, CAL. CAL reveals whether you are losing the connective tissue foundation that holds the tooth. For example, a 2 millimeter probing depth with 2 millimeters of recession equals 4 millimeters of attachment loss, which is different from a 4 millimeter pocket with no recession. Same number, different biology. Bleeding on probing tells you about ongoing inflammation. Suppuration, mobility grades, and any furcation involvement on molars fill out the risk profile.

A winnipeg dentist will often record a plaque score as well, such as an O’Leary index, to quantify how much biofilm clings to the teeth and where it concentrates. Numbers turn a vague sense of “I brush well” into a map that shows the truth.

Photos, scans, and the value of evidence you can see

Recession is easier to discuss with images. Intraoral photographs, taken with cheek retractors and a few mirrors, capture the gingival margins clearly. Baseline photos are priceless, because the human eye normalizes slowly shifting lines. A year later, you can compare pixel to pixel.

Many clinics now use intraoral scanners to create 3D models of the teeth and gums. While scanners are stellar for bite alignment and restorations, they also capture the gingival margin. Side-by-side scans six months apart can show a half-millimeter retreat that would have been missed in conversation.

Radiographs, and when a CBCT is worth the radiation

Bitewings reveal interproximal bone levels and root caries risk. Periapicals help when a single tooth is a problem, especially if a root looks prominent under a thin gum. Radiographs do not show soft tissue, so you never diagnose recession from an x-ray, but you can spot the bone architecture that supports the gum.

Cone beam CT is not routine for recession. It enters the picture for complex cases, particularly when orthodontic movement is planned or if fenestrations and dehiscences are suspected in the facial plate. You do not want to move a tooth further outside the bony envelope when the gum is already protesting.

The occlusion check: your bite leaves fingerprints

Teeth that carry too much load rock over time, sometimes microscopically, and the gumline tells the tale. The dentist watches how your teeth meet, asks you to slide forward and side to side, and looks for shiny wear facets. Abfraction lesions near the gumline often pair with heavy lateral contacts. An occlusal guard is not a cure for recession, but it reduces one source of trauma while the gums heal and adapt.

Frena, piercings, and other local troublemakers

That little string of tissue between your lip and gum, the frenum, should be a quiet background character. Sometimes it inserts sevenoaksdentalcentre.com too close to the margin and tugs at the tissue with every word. Add a narrow band of keratinized tissue and a toothbrush, and recession follows. A labial frenum on a lower incisor can be a small thing with outsized influence.

Oral piercings rub. A metal ball clicking against the lower incisors all day does not get filed under harmless. Many a dentist in winnipeg has a story of a tongue piercing retired for the sake of four unhappy front teeth.

Sorting causes: periodontal disease, mechanical trauma, or both

Not all recession is inflamed. This is a crucial distinction. Some defects have no redness, no plaque, and no bleeding, just thin tissue and exposed root. These cases lean toward mechanical etiology: overzealous brushing, orthodontic movement beyond the bony envelope, or a tight frenum.

Other sites bleed upon gentle probing, carry plaque, and sit next to 5 to 6 millimeter pockets. Here, recession is a face of periodontitis. Treatment priorities shift. You address inflammation first with meticulous debridement and home care instruction, then reassess the margins.

Certain desquamative conditions mimic or complicate recession. Lichen planus, mucous membrane pemphigoid, or a reaction to cinnamon oil can make the gums tender and shiny, which changes how and where the patient brushes. A good history and, if needed, a biopsy through a referral answer these edge cases.

Classification that guides decisions

Classification systems translate millimeters into prognosis. The older Miller classification tied recession to interdental bone loss and the mucogingival junction. Many clinicians now prefer Cairo’s system, which frames it around interproximal attachment loss measured with CAL.

- RT1: recession with no interproximal attachment loss. These cases have excellent potential for full coverage with grafting if indicated.
- RT2: recession with interproximal loss that is less than or equal to the facial loss. Coverage potential is partial to substantial.
- RT3: recession with interproximal loss greater than the facial loss. Coverage is limited.

Why bother with labels? Because they align expectations. If a patient expects a magic graft to return the gumline to its teenage position while proximal bone is gone, you need a common language to explain biology and limits.

Two true Winnipeg stories, anonymized and instructive

A mid-30s hockey coach came in every March with the same complaint: the lower left canine zapped him whenever he drank anything colder than room temperature. Clinical charting showed 2 millimeters of recession, no bleeding, a thin biotype, and a narrow band of keratinized tissue. There was a gleaming toothbrush abrasion notch. His brushing demo was a vigorous saw motion he had perfected over years. We switched him to a soft powered brush with a pressure sensor, taught a modified stillman technique angled partly onto the gum, applied a varnish for sensitivity, and added a custom night guard. Six months later, photos showed no further recession. Two years later, still stable. He decided to skip grafting unless the site changed. Sometimes the fix is behavior and patience.

A retiree in River Heights came in with generalized sensitivity and one front tooth that looked longer every spring. We found 4 to 5 millimeter pockets on multiple teeth, bleeding on probing at 30 percent of sites, and 3 millimeters of recession on the upper right canine. Radiographs showed moderate interproximal bone loss. Here the recession was part of periodontitis. We scheduled quadrant scaling with local anesthesia, calibrated home care to a small head brush and tapered interdental brushes that fit her spacing, and checked her HbA1c with permission from her physician because her diabetes had been wobbly. Three months after debridement, bleeding dropped to 10 percent and the tissue tone improved. The canine recession was remeasured and found stable. We referred her to a periodontist to discuss grafting options, with the understanding that perfection was unlikely, but root coverage and thicker tissue could reduce sensitivity and future risk.

What to expect during a recession-focused visit

For many new patients, a dentist in winnipeg will fold this evaluation into a comprehensive exam. If you are booking specifically for recession, here is the short version of the flow.

- A focused history that covers habits, symptoms, medical context, and past dental work.
- A thorough periodontal charting with six sites per tooth, plus mobility and furcations when relevant.
- High quality photos and selective radiographs to document baselines and rule out hidden caries or bone issues.
- An occlusion review with attention to grinding patterns and abfraction clues.
- A discussion that blends numbers, images, and realistic options, from habits to grafting.

A good clinician never jumps straight to surgery, and never dismisses surgery when it is the right choice. Judgment lives in the middle.

When recession is sensitive but stable

Sensitivity alone does not equal progression. If the tissue is pink, firm, and non-bleeding, and photos show no change across six to twelve months, the dentist may stick to conservative measures.

Desensitizing toothpaste with 5 percent potassium nitrate or arginine can calm nerves. A high fluoride varnish or silver diamine fluoride spot application reduces root caries risk and often cuts sensitivity. Recontouring a sharp composite edge or smoothing a toothbrush notch can help. Correcting brushing technique matters more than any gadget. The instructions are simple and maddening: slight angle toward the gum, gentle pressure you could use on a ripe peach, short strokes or small circles, two minutes, twice daily. Most people need coaching, not a lecture.

When thin tissue needs a bodyguard

If the biotype is very thin and the band of keratinized tissue is narrow or absent, recession tends to march on. Here, the evaluation turns to grafting candidacy. The periodontist or a general dentist with advanced training evaluates tissue thickness with a probe transparency test, confirms Cairo classification, and maps local anatomy. Options now include connective tissue grafts from the palate, using a tunneling or coronally advanced flap approach, or a collagen matrix substitute that spares the palate in selected cases. Full root coverage is not guaranteed, but a thicker, tougher gum often is. That thickness is a bodyguard for the long term.

Patients always ask, how bad does it hurt? Experience varies. Palatal donor sites are the sore part. A collagen matrix cuts that down, though sometimes at the cost of slightly less predictability. Most people manage with over the counter analgesics and a week of chewing on the other side. A winnipeg dentist will time grafting around your calendar, not in the middle of playoff season if you coach four nights a week.

Orthodontics and the bony envelope lesson

Teeth like to live inside bone. Move them too far outward, and the bone thins, then the gum thins, then it recedes. This does not mean orthodontics causes recession by default, but movement plans must respect the bony envelope. If you are straightening in adulthood and your lower incisors are already flirting with thin tissue, your orthodontist and your general dentist should talk. Sometimes the right sequence is graft first, move second. Occasionally it is the other way around.

For clear aligner fans, attachments and trays make cleaning trickier, which boosts inflammation if you are not diligent. Recession plus gingivitis is a predictable recipe for frustration. A simple rule: if you cannot clean it easily, you will not clean it well. Plan for more frequent hygiene visits during orthodontics.

The bruxer’s paradox

People who grind often brush harder. It is a personality adjacency, not a law. If you clench at night, a properly fitted guard relieves lateral forces. That alone does not stop recession, but it reduces one line of attack on the necks of the teeth. Pair it with a softer brush and, if needed, a periodic review of stress management. Your gums will not thank you in a note, but the probe readings will.

Smoking, vaping, and the quiet bleed

Nicotine chokes microcirculation. Smokers bleed less even when inflamed, which tricks the eye. A site might look calmer than it is. Vaping, depending on the fluid cocktail, can dry the mucosa and irritate tissue. During an exam, a dentist winnipeg patients see regularly will fold tobacco use into risk calculus and adjust expectations accordingly. Grafts do better in non-smokers, plain and simple.

Costs, coverage, and the honest conversation

Evaluating recession takes time, and so does treating it. In Manitoba, fee guides help set expectations, but coverage varies widely. A winnipeg dentist typically provides a written plan with codes and fees you can submit to your insurer. Even if grafting is on the horizon, the first investments are usually the cheapest and most powerful: coaching on home care, a soft brush or a powered one with a pressure sensor, a custom guard if you grind, and targeted cleaning visits. If surgery follows, you will feel more confident because you already saw stability improve.

How to keep score at home between visits

You are part of the evaluation team whether you like it or not. Watching your own gumline is less about obsession, more about pattern recognition.

- Once every month or two, take a clear smartphone photo of the area in good light with your lip gently lifted. Compare to the last one rather than to memory.
- Pay attention to sensitivity triggers. If cold pain increases over weeks despite using a sensitivity paste, mention it.
- Notice bleeding points when you floss. A spot that bleeds every time deserves a look.
- Track any change in brushing tools or techniques. If you switch brushes and the site worsens, tell your dentist.
- If you start or change medications, especially those affecting hormones or immune response, keep the dental team in the loop.

You do not need calipers in your bathroom drawer. You do need the curiosity to notice a trend.

Red flags that speed up the timeline

Even the calmest watch-and-wait plan can pivot. Rapidly increasing sensitivity, visible root caries, a frenum that whitens and pulls the margin when you smile, mobility you can feel with your tongue, or a site that starts bleeding after years of quiet, all of these push a dentist in winnipeg to move from conservative to interventional. It might be time for a graft consult or a bite adjustment. It might be as simple as repairing a notched margin that catches the brush and magnifies trauma.

A dentist’s calculus: sequence, not just selection

The biggest difference between a quick look and a professional evaluation is sequencing. The order matters.

First, reduce active inflammation and mechanical trauma. Second, stabilize habits with coaching and tools that fit your life. Third, document stability with photos and charting. Fourth, if the tissue is thin or defects are deep, plan for grafting with realistic goals. Fifth, maintain with regular hygiene that respects your risk level, which could be every three months for a while, then back to six.

This sequence is not dogma, it is rhythm. When you follow it, results feel less like a lucky break and more like a well played shift.

The short version, if you only remember one thing

Gum recession is not a single problem with a single fix. It is a pattern with multiple inputs, and the way a winnipeg dentist evaluates it reflects that complexity. History, measurements, images, bite, tissue type, and your daily habits all get weighed. The magic is not in a fancy probe or a perfect graft, it is in the way information gets put together, communicated, and acted on over time.

If your gumline is making you nervous, book the visit. Bring your questions and your toothbrush, and be ready to talk about how you use it. The millimeters will make sense by the time you leave, and that small sense of control is worth almost as much as the numbers themselves.

Seven Oaks Dental Centre
< a href="https://www.sevenoaksdentalcentre.com">https://www.sevenoaksdentalcentre.com
Address: 2110 Main St, Winnipeg, MB R2V 2C2

Phone: +12043344388