

Most people do not think about dentistry in categories. They think in moments. A tooth starts to ache during dinner. A child chips an incisor on the playground. A hygienist mentions early gum inflammation at a routine visit. A dentist spots a cavity on a bitewing X-ray that the patient could not feel at all. General dentistry sits right in the middle of those ordinary moments. It is the part of dental care that handles prevention, diagnosis, maintenance, and many of the treatments that keep small problems from becoming expensive, painful ones.

When patients ask what counts as a “common” treatment, they are usually asking two things at once. First, what procedures are performed most often in a general dental office? Second, which of those procedures are most likely to affect me or my family? The answer is broader than many people expect. General dentistry is not limited to cleanings and fillings, although those are certainly central. It also includes exams, X-rays, fluoride treatments, sealants, periodontal care, crowns, simple extractions, and treatment for worn or damaged teeth. In many practices, it even overlaps with cosmetic, emergency, and restorative care.

The common thread is practical care. General dentistry focuses on keeping the mouth healthy, functional, and stable over time. That often means treating disease early, watching areas that are not yet severe enough to treat, and helping patients make decisions that balance cost, longevity, comfort, and appearance.

Routine exams and professional cleanings

If one treatment defines general dentistry, it is the routine checkup paired with a professional cleaning. This sounds simple, but it is the foundation of nearly everything else. A dental exam is not just a quick look at the teeth. A thorough visit usually includes an evaluation of the gums, tongue, cheeks, bite, existing dental work, and signs of wear or grinding. Dentists also check for changes in soft tissues, which is one reason regular visits matter even for people who rarely get cavities.

The cleaning itself, often performed by a dental hygienist, removes plaque and tartar that brushing and flossing cannot fully manage at home. Plaque is soft and can usually be disrupted with good home care. Tartar, or calculus, hardens on the teeth and must be removed with professional instruments. Once tartar builds up around the gumline, it creates a rough surface that attracts more plaque, which makes inflammation harder to control.

A common misconception is that if teeth look white and feel smooth, there is nothing to worry about. In practice, the earliest gum disease often causes little pain. Mild bleeding during flossing is one of the most overlooked warning signs in dentistry. Many patients assume bleeding means they should floss less. Usually the opposite is true, though technique matters. A professional cleaning resets the environment, and consistent home care helps maintain it.

The interval between visits varies. Six months is common, but it is not universal. Someone with a history of gum disease, heavy tartar buildup, dry mouth, or frequent decay may benefit from more frequent maintenance, often every three or four months. A low-risk adult with excellent home care and little dental history may not need that pace. Good general dentistry is individualized, not automatic.

Dental X-rays and diagnostic imaging

X-rays are another common part of general dental care, and patients often underestimate how much they reveal. Many cavities begin between teeth where they are not visible to the eye. Bone loss from gum disease can also progress silently before symptoms become obvious. A cracked filling, an infection at the root tip, or an unerupted tooth may only show up on imaging.

Bitewing X-rays are among the most frequently taken images in general dentistry because they help detect decay between back teeth and show bone levels around those teeth. Periapical images give a more complete view of the entire tooth and root. Panoramic X-rays are less routine for every recall visit, but they can be useful for seeing the broader picture, including wisdom teeth, jaw structures, and some pathology. Many offices now use digital radiography, which reduces radiation compared with older film systems and makes images available immediately.

The value of X-rays is timing. It is much easier to repair a small cavity than to save a tooth that has developed a deep infection because decay went unnoticed for too long. Patients who **General Dentistry** want to skip imaging often do so because nothing hurts. Unfortunately, discomfort is a poor screening tool for early dental disease. Many serious problems become painful only after they are advanced.

Fillings for cavities and small fractures

Tooth-colored fillings remain one of the most common treatments in general dentistry. They are used to repair cavities, replace broken portions of teeth, and sometimes remove and update older restorations that have worn down or developed leakage. Composite resin is now the standard material in many offices because it bonds to tooth structure and blends well with natural enamel.

From the patient's perspective, a filling can seem minor. Clinically, the details matter. A tiny cavity confined to enamel is very different from a broad cavity that extends deep into dentin near the nerve. The larger the decay, the more difficult it is to preserve strength and avoid future complications. This is one reason dentists emphasize routine exams. They are not trying to "find work." They are trying to catch restorations while they are still straightforward.

There is also judgment involved in deciding when to treat. Not every stained groove is decay. Not every shadow on an X-ray needs immediate drilling. In experienced hands, diagnosis includes watchful monitoring when appropriate. Some early lesions can be managed with fluoride, improved hygiene, and diet changes, especially if the outer tooth surface is still intact. Once a cavity has clearly broken through and softened the tooth, a filling is usually the practical next step.

Patients often ask how long a filling lasts. There is no honest single number. A small filling in a low-stress area may last many years. A large filling in a patient who clenches at night may fail sooner. Diet, home care, bite forces, and the size of the restoration all matter. The best way to make a filling last is to need the smallest filling possible in the first place.

Fluoride treatments and sealants

Not every common dental treatment involves repairing damage. Some of the most useful services are preventive. Fluoride treatments are especially common in children, but adults can benefit too, particularly those with dry mouth, gum recession, orthodontic appliances, high cavity risk, or a history of repeated decay.

Fluoride strengthens enamel and helps teeth resist acid attacks from plaque bacteria and diet. In an office setting, it is usually applied as a varnish, gel, or foam after a cleaning. The process is quick, but its value can be significant in the right patient. I have seen adults with medication-related dry mouth go from getting frequent root cavities to stabilizing well once fluoride, saliva support, and home care were taken seriously.

Sealants are another preventive [General Dentistry](#) staple, mostly for children and teenagers but sometimes useful for adults with deep grooves in their molars. The chewing surfaces of molars have pits and fissures that are ideal hiding places for plaque and food debris. A sealant is a thin protective coating placed over those grooves to reduce the risk of decay. When placed well and monitored over time, sealants can be highly effective.

These treatments do not replace brushing, flossing, or dietary discipline. They support them. General dentistry works best when prevention is layered, not when any one product or procedure is expected to do all the work.

Gum disease treatment beyond the routine cleaning

Patients often use the phrase “deep cleaning” casually, but periodontal treatment is not just a more intense version of a regular prophylaxis. It addresses disease under the gumline, where bacteria and calculus trigger inflammation that can damage supporting bone. In early stages, gum disease may present as bleeding, puffiness, or bad breath. Later on, it can lead to pocketing, gum recession, mobility, and tooth loss.

Scaling and root planing is one of the most common periodontal procedures in general dentistry. It involves cleaning below the gumline to remove deposits from root surfaces and reduce bacterial load. Depending on the extent of the disease, local anesthetic may be used for comfort, and treatment may be completed by sections of the mouth. Afterward, patients usually enter a periodontal maintenance schedule rather than simply going back to standard cleanings twice a year.

This distinction matters. A routine cleaning is for a generally healthy mouth or one with mild gingivitis. Periodontal maintenance is for someone with a history of periodontal disease that needs closer control. The bone lost to periodontitis does not simply grow back in most everyday cases, so long-term management is essential.

One of the most frustrating realities in dentistry is that gum disease can advance in people who think they are doing everything right. Sometimes brushing technique misses the gumline. Sometimes flossing is inconsistent. Sometimes smoking, diabetes, genetics, or dry mouth complicates the picture. Good general dentistry is careful not to blame patients simplistically. It identifies risk factors, explains what can be changed, and sets realistic expectations.

Crowns for weakened or heavily restored teeth

When a tooth has lost too much structure for a filling to hold up predictably, a crown often becomes the treatment of choice. Crowns cover and protect the visible part of the tooth, restoring strength, shape, and function. In general dentistry, crowns are commonly recommended after a large cavity, a fracture, root canal treatment, or repeated replacement of older restorations.

The decision between a large filling and a crown is one of the most common judgment calls in practice. Patients sometimes prefer the less expensive option in the short term, which is understandable. But when a tooth has thin remaining walls, a very large filling may act more like a wedge than a support. Under chewing pressure, the tooth can crack. If the crack stays above the gumline, the tooth may still be savable with a crown. If it extends deeper, the tooth may be lost.

Modern crowns can be made from several materials, including all-ceramic and porcelain-fused-to-metal options. The best choice depends on where the tooth is located, how hard the patient bites, and aesthetic priorities. A crown on a front tooth has different demands than one on a back molar in a patient who clenches heavily.

Patients often ask whether getting a crown means the tooth was neglected. Not necessarily. Some teeth simply reach the end of what a filling can reasonably support. A person may have had a large filling placed years ago, and the crown is the next sensible step when that restoration wears out or the tooth structure weakens. General dentistry often involves extending the useful life of a tooth through stages of care.

Root canal treatment when the nerve is involved

Although some root canal therapy is referred to endodontists, many general dentists perform it routinely on selected teeth. This treatment becomes necessary when the pulp, the inner nerve and blood supply of the tooth, becomes inflamed or infected. The causes are familiar: deep decay, trauma, cracks, or repeated procedures on the same tooth.

The symptoms vary more than most people expect. Some patients have severe throbbing pain, sensitivity to biting, or swelling. Others have a dead tooth with little pain at all, discovered only when an X-ray shows infection at the root tip. That surprise is common. Teeth do not always read the textbook.

During root canal treatment, the dentist removes the infected pulp tissue, cleans and shapes the canals, disinfects the space, and seals it. In many cases, the tooth then needs a crown because a tooth that has had root canal therapy is often more brittle and structurally compromised than before. Saving the tooth is usually the goal because maintaining a natural tooth, when feasible, helps preserve biting function and reduces the need for replacement options.

Root canals suffer from an outdated reputation. The procedure itself is usually not the ordeal patients fear. The real problem is waiting too long while the tooth is already badly infected. Prompt treatment generally means a smoother experience and a better prognosis.

Extractions and when removing a tooth is the right call

General dentistry is centered on saving teeth whenever possible, but not every tooth can or should be saved. Simple extractions remain common, especially for teeth that are severely decayed, broken beyond repair, advanced in gum disease, or causing crowding or infection. Some general dentists also remove certain wisdom teeth, though more complex surgical cases are often referred out.

No experienced dentist recommends extraction lightly. Once a tooth is gone, the consequences ripple outward. Neighboring teeth can drift, opposing teeth can over-erupt, chewing patterns can change, and bone in the area gradually resorbs. That is why dentists often discuss replacement options such as implants, bridges, or partial dentures after extraction. The best decision depends on age, budget, bone support, health history, and how important that tooth is to the patient's bite.

There are edge cases where extraction is the better decision even if a heroic save is technically possible. A tooth with a poor crack pattern, limited remaining structure, heavy bite stress, and a guarded long-term outlook may consume a great deal of money and time without giving the patient reliable service. One hallmark of strong general dentistry is candor. Saving a tooth should be meaningful, not symbolic.

Treatment for tooth wear, grinding, and sensitivity

Not all common dental treatment revolves around decay. Tooth wear is increasingly common, and it shows up in patients of every age. Some grind at night. Some clench during the day without realizing it. Others sip acidic drinks all afternoon, creating chemical wear that softens enamel over time. Recession can expose root surfaces, leading to sensitivity and a higher risk of root decay.

General dentists manage these issues in several ways. Sometimes the solution is a night guard to protect against grinding forces. Sometimes it is bonding to repair worn edges. Sometimes it involves fluoride, desensitizing agents, or changes in brushing technique. Hard scrubbing with a medium or firm brush can do real damage over the years, especially near the gumline. A soft brush used well is usually the better tool.

This category of care often requires patience because the treatment is not always a single appointment fix. A patient with cold sensitivity might need an adjustment in home products, diet, brushing habits, and bite

protection before symptoms settle. The best results usually come when the dentist connects the dots between symptoms and habits, rather than treating sensitivity as an isolated complaint.

Care for children and family patients

A great deal of General Dentistry happens in family settings, where care needs shift by age. For children, common treatments include exams, cleanings, fluoride, sealants, monitoring eruption patterns, and treating cavities in both baby and permanent teeth. Early visits also shape comfort. A child who learns that dental appointments are predictable and nonthreatening often becomes an adult who seeks care earlier and more consistently.

For teenagers, sports guards, sealants, orthodontic referrals, and management of diet-related decay are common themes. Sugary drinks, frequent snacking, and inconsistent brushing can undo a lot of good intentions. For adults, the pattern often changes to maintenance of older fillings, crowns, gum health, and wear from stress or aging. For older adults, dry mouth, recession, root caries, and management of complex restorative histories become especially important.

The treatment names may sound familiar across these life stages, but the context changes. A small cavity in a six-year-old first molar is not the same conversation as a failing large restoration in a sixty-year-old molar with a crack line. General dentistry is common precisely because it follows patients through those transitions.

What determines which treatment you actually need

Two patients can sit in the same waiting room and receive completely different recommendations, even if both say, "Nothing hurts." That is normal. Dental treatment is shaped by several practical factors:

1. Current disease activity, such as new cavities, gum inflammation, or a cracked tooth.
2. Risk level, including dry mouth, diet, home care, smoking, and previous dental history.
3. Structural condition of the tooth, especially how much healthy tooth remains.
4. Bite forces and habits like clenching, grinding, nail biting, or chewing ice.
5. Long-term goals, budget, and whether the patient wants the most conservative or most durable option.

That final point matters more than people realize. Good dentistry is not just about diagnosing correctly. It is also about matching treatment to the patient's reality. A crown may be the ideal restoration on paper, but a well-planned interim filling may be the practical step if finances are tight and the tooth can be stabilized safely. On the other hand, repeatedly patching a failing tooth can cost more in the long run than addressing it definitively.

The treatments patients end up needing most often

If you strip general dental care down to what most patients are most likely to encounter over time, the usual sequence is fairly predictable. People start with preventive care, then receive repair work if disease or wear develops, and move into more protective or restorative procedures as teeth age.

In everyday practice, the most common treatments are routine exams and cleanings, X-rays, fillings, fluoride or sealants for prevention, gum disease treatment when needed, crowns for weakened teeth, and occasional root canals or extractions when problems are advanced. None of these exists in isolation. A cleaning may uncover gum disease. An X-ray may reveal a cavity that only needs a small filling because it was found early. A large filling may preserve a tooth for years before a crown becomes the wiser choice.

That is the practical value of General Dentistry. It is not glamorous, and it does not need to be. Its purpose is to keep ordinary dental problems ordinary. The earlier they are seen, the simpler the treatment tends to be. The longer they are ignored, the narrower the options become. For most patients, the most common dental treatments are also the most preventable, which is exactly why regular care matters so much.

Aspenwood Dental Associates and Colorado Dental Implant Center

Address: 2900 S Peoria St Ste C, Aurora, CO 80014

Phone number: +13037314037

FAQ About General Dentistry Aurora

What is meant by general dentistry?

General dentistry refers to the primary, foundational tier of oral healthcare, focused on the prevention, diagnosis, and treatment of conditions affecting the teeth, gums, and jaw. General dentists serve as a patient's main, long-term dental care provider—much like a primary care physician.

What is general dentistry and orthodontics?

General dentistry and orthodontics are two specialized branches of dental care. General dentistry serves as your primary care for overall oral health, focusing on routine cleanings, fillings, and disease prevention. Orthodontics is a specialized field focused entirely on diagnosing and correcting misaligned teeth and jaw structures using braces or clear aligners.

What are type 3 dental services?

Type 3 dental services typically include major restorative treatments that repair or replace damaged or missing teeth. These services are more complex and costly than preventive or basic dental care. Common examples of type 3 dental services include: Dental crowns.